

Comparative studies on anti-avian sera conjugated with fluorescein isothiocyanate

M. H. Khodeir^{1*}, Ghada M. El-Sadek², M. M. El-Safty², Elham A. El-Ibiary²

¹*Veterinary Serum and Vaccine Research Institute, Cairo and* ²*Central Laboratory for Quality Control of Veterinary Biologics, Cairo, Egypt.*

Rabbit antisera were successfully prepared against chickens; turkey; ducks; geese; pigeons and quails as antispecies and conjugated with fluorescein isothiocyanate (FITC). The ability to use these antisera as homologous and heterologous antispecies was studied through the application of indirect fluorescent antibody technique (FAT) using specific antigens and antibodies of Newcastle; infectious bursal disease virus; duck plague virus; duck hepatitis virus; fowl cholera and pigeon paramyxo virus. The results showed that Titration of the prepared anti avian sera conjugated with FITC induced strong positive FAT reactions up to dilutions of 1:10000; 1:1000; 1: 100000; 1: 1000; 1: 100000 and 1:1000 for anti- chicken; anti-Turkey; ant-duck; ant-geese; anti-pigeon and anti-quail sera respectively. It was found that homologous species anti-sera showed strong positive FAT reaction (green apple fluorescing) scored as 4+ while heterologous species anti-sera showed less degree of positive reactions ranged from 3+ to ± reaction showing that the usefulness of anti-sera as reagents in serological techniques is limited by the homogeneity and heterogeneity of such anti-sera. So, the present preparations could be used to low extent as heterologous anti-avian sera.

Specific antisera to individual immunoglobulin classes are valuable reagents in many immunological techniques such as indirect immunofluorescent technique (Kincade and Cooper, 1971-1973; Lawrence *et al.*, 1979). Specific, rapid and sensitive serological tests like fluorescent antibody technique (FAT) are required for accurate diagnosis of infectious agents to reach to well applicable control measures. FAT depends mainly on antigen-antibody reaction in the presence of a fluorescent dye (fluorescein isothiocyanate "FITC") which irradiates with ultraviolet light emitting apple green fluorescence (Tizard, 1996). The technique depends on the presence of specific antiserum conjugated with FITC. FAT has gained wide acceptance in virology serving for many purposes such as detection of cellular localization of viral antigens (Watson and Coons, 1954; Spendlove *et al.*, 1963); establishment of temporal sequence of antigens after infection appearance (Breitenfeld and Schafer, 1957); and detection of virus infected cells which fail to produce viral progeny and still synthesize virus coded antigens which continue to harbor viral genetic material (Pope and Rowe, 1964; Tevethia *et al.*, 1965).

The degree of specificity of the antiserum may have a considerable influence on the results.

Commercially available antisera for use in avian serological assays are generally lacking in specificity (Nadaphan *et al.*, 1982). In addition, the use of heterogenous antibodies provides less potential for the immobilization to impact the antigen binding efficiency (Tumpey *et al.*, 2005).

The present work was designed to prepare and compare between different specific anti-avian sera (chicken; turkey; ducks; geese; pigeons and quails) conjugated with fluorescein isothiocyanate concerning their specificity and the ability to use any of them instead of the other as heterologous agent in the absence of the homologous one.

Materials and methods

Birds. Five 2 months old of each of chickens; turkey; ducks; geese; pigeons and quails were used to collect blood and sera to be used for preparation of anti-species sera in rabbits. All birds were clinically healthy and free from Newcastle; infectious bursa disease; duck plague; duck hepatitis; fowl cholera and pigeon paramyxo antibodies and housed under hygienic measures in separated isolates receiving balanced ration and adequate water.

Rabbits. Thirty five healthy Bosket rabbits of average body weight 3 kg were used for preparation of different anti-avian sera whereas each antiserum was prepared in five rabbits leaving five rabbits as control. Each rabbit group was housed separately in hygienic isolates.

Preparation of species anti-sera. It was carried

* Corresponding author. Tel.: +202 3424406;
 Fax: +202 3428321
 E-mail address: svri@idsc.gov.eg
 (M. H. Khodeir).




Fig. (1-6): 4+; 3+; 2+; +; ± and -ve FAT are shown in fig. 1,2,3,4,5 and 6 respectively

out following the procedure described by (Deutsch, 1967).

Precipitation of the immunoglobulin of the prepared antisera. It was carried out using ammonium sulphate according to (Narin and Marrack, 1964). The globulin content was estimated according to (Weichselbaum, 1946) and adjusted to be 20mg/ml using phosphate buffer saline of pH 7.2.

Chemicals used for conjugation of the prepared species antisera with Fluorescein isothiocyanate (C₄ H₁₁ NO₅ S) E. (C₄ H₁₁ NO₅ S) E was supplied by Merck, Darmstadt for Microscopy (M.Gew.389.39).

Conjugation of the prepared species antisera with Fluorescein isothiocyanate. It was done according to the method of (Narin, 1969).

Antigens and antibodies. Newcastle; infectious bursa disease; duck plague; duck hepatitis; fowl cholera and pigeon paramyxo antigens and antibodies were supplied by the Central Laboratory for Quality Control of Veterinary Biologics, Abassia; Cairo.

Reference anti-chicken conjugate. Fluorescein-labeled affinity purified antibody to chicken IgG (H+L) produced in goat, catalog No.02-24-06 was supplied by KPL Europe. It was used as a

reference conjugate to evaluate the prepared anti-avian sera conjugated with FITC. It was used at a dilution of 1:100 according to the manufacturer directions.

Indirect fluorescent antibody technique (FAT). It was carried out on Newcastle; infectious bursa disease; duck plague; duck hepatitis; fowl cholera and pigeon paramyxo antigens and antibodies using the prepared species antisera conjugated with fluorescein isothiocyanate in homologous and heterologous manners. The technique was carried out according to (Soliman *et al.*, 1989).

Results and Discussion

Usually and especially in recent works, there is an increased demand to obtain rapid, sensitive and accurate diagnosis of infectious diseases especially those occur in the form of outbreaks. One of the most reliable techniques is the fluorescent antibody technique (FAT) which provides such requirements. Indirect FAT depends mainly on the presence of specific anti-specie serum conjugated with fluorescein isothiocyanate (Brian and Hillor, 1996; Hala *et al.*, 2003; Hanan *et al.*, 2003; Noura *et al.*, 2009).

The present work was mainly directed to prepare specific antisera against some avian species conjugated with FITC to investigate to any extent such antisera could be used instead of each other.

All prepared antisera showed higher levels of total protein level than in normal rabbit sera due to the increase in globulin levels from which species antibodies are formed (Table-1). These results are in accordance with those of Kataria and Sharma, (1993); Hala *et al.*, (2003); Hanan *et al.*, (2003).

Concerning titration of the prepared anti-avian sera conjugated with FITC (Table-2), it was found that these preparations induced strong positive FAT reactions up to dilutions of 1:10000; 1:1000; 1: 100000; 1: 1000; 1: 100000 and 1:1000 for anti- chicken; anti-Turkey; ant-duck; ant-goose; anti-pigeon and anti-quill sera respectively. These values appear to be higher than that of the imported references anti-chicken conjugate which induced positive reaction only up to a dilution of 1:100 which was prepared according to the manufacturer directions. So, the present preparation could be considered preferable specific products than the non specific imported one whereas they provide larger amounts of the conjugate (when diluted on use)

Table (1): Serum proteins in the prepared rabbit anti-avian sera.

Tested antiserum	Total protein (g/dl)	Albumin (g/dl)	Globulin (g/dl)
Anti-chicken	5.57	2.35	3.22
Anti-Turkey	5.62	2.04	3.58
Anti-duck	5.79	2.37	3.42
Anti-goose	5.40	2.03	3.37
Anti-pigeon	5.12	2.25	2.87
Anti-quill	5.45	1.85	3.60
Normal rabbit serum	4.26	2.03	2.23

Table (2): Titer of FITC conjugated anti-avian sera.

Conjugate dilution	Tested conjugated antisera					
	Anti-chicken	Anti-Turkey	Anti-duck	Anti-goose	Anti-pigeon	Anti-quill
Undiluted	+	+	+	+	+	+
1:10	+	+	+	+	+	+
1:100	+	+	+	+	+	+
1:1000	+	+	+	+	+	+
1:10000	+	-	+	-	+	-
1:100000	-	-	+	-	+	-
1:1000000	-	-	-	-	-	-

NB: The reference antichickens conjugate was diluted as 1:100 according to the manufacturer directions.

Table (3): Positively reacted FITC conjugated homologous and heterologous anti-avian sera.

Tested anti-sera	Indirect FAT using					
	Chicken serum	Turkey serum	Duck serum	Goose serum	Pigeon serum	Quill serum
Anti-chicken	4+	3+	2+	2±	2+	+
Anti-Turkey	3±	4+	+	±	+	-
Anti-duck	3+	2+	4+	4±	±	-
Anti-goose	2+	+	4+	4+	±2	±
Anti-pigeon	3±	2±	+	2±	4+	3+
Anti-quill	2±	+	±	+	4±	4+

saving cost. Similar findings were recorded by (Hala *et al.*, 2003; Hanan *et al.*, 2003; Abd El-Wanis and Khodeir, 2004; Hussein *et al.*, 2006;

Regarding the use of homologous and heterologous species anti-sera; it was found that homologous species antisera showed strong positive FAT reaction (green apple fluorescence) scored as 4+ while heterologous species anti-sera showed less degree of positive reactions ranged from 3+ to ± reaction as demonstrated in table (3). Such scores of positive FAT reactions were demonstrated in (Fig. 1-6). These findings confirm previous records of (MRC Working Shop, 1971; Hala *et al.*, 2003; Hanan *et al.*, 2003) showing that the usefulness of anti-sera as reagents in serological techniques is limited by the homogeneity and heterogeneity of such antisera.

It could be concluded that the prepared anti-avian sera conjugated with FITC are of good quality and could be used for detection of Newcastle; infectious bursa disease; duck

Noura *et al.*, 2009) who prepared different local antisera conjugated with FITC of high quality saving time and cost.

plague; duck hepatitis; fowl cholera and pigeon paramyxo antigens and antibodies with high specificity and sensitivity saving the long time and high cost required for importation of such conjugates. In addition, the present preparations could be used to low extent as heterologous anti-avian sera.

References

- Abd El-Wanis, N. A. and Khodeir, M. H. (2004): Conjugation of immunoglobulins prepared against infectious bursal disease virus in different hosts with fluorescein isothiocyanate. *Egypt. J. Agr. Res.*, 82(4): 1881-1891.
- Benedict, A. A. (1967): *Methods in immunology and immunochemistry*. Academic press, London and New York. P.294
- Breitenfeld, P. and Schafer, W. (1957): Sequence of different antigens. *Virology*, 4: 328-333.
- Brian, W. J. and Hillor, O. (1996): *Virology methods manual*. P.318. Academic press, London, San Diego, New York, Sydney, Tokyo.

- Deutsch, H. F. (1967):** Methods in immunology and immunochemistry. Academic press, London and New York. P.315
- Hala, A. F.; Khodeir, M. H. and Hussein, A. Z. (2003):** Preparation and evaluation of rabbit antigoat and antisheep serum conjugated with fluorecein isothiocyanate. Beni-Suef Vet.Med.J. Vol.XIII, No.(1): 125-133.
- Hanan, M. I.; Zeinab, M. S.; Elham, A. E. and Khodeir, M. H. (2003):** Preparation and evaluation of antiduck and antiturkey sera conjugated with fluorecein isothiocyanate for serological use. Beni-Suef Vet.Med.J. Vol.XIII, No. (1): 193-199.
- Hussein, A. S.; Khodeir, M. H.; Fathia S. and Hussein, A. Z. (2006):** Conjugation of Clostridium chauvoei and Clostridium septicum hyper immune sera with fluorescein isothiocyanate for serological uses. Minufiya vet.J.Vol.4 (2): 313-317.
- Kataria, A. K. and Sharma, K. N. (1993):** Serum electrophoretic studies on sheep vaccinated with live sheep pox virus (Rumanian strain). Ind.Vet.J.,50: 387-393.
- Kincade, P. W. and Cooper, M. D. (1971):** Momnospecific antisera. J. Immunol., 106: 371-376.
- Kincade, P. W. and Cooper, M. D. (1973):** Immunoglobulin classes. Cellular immunology, 8:93-98.
- Lawrence, E. C.; Arnaud-Battexdier, F.; Koski, I. R.; Dooley, N. Z.; Mughmore, A. V. and Blaese, R. M. (1979):** Immunological techniques (FAT). J. Immunol., 123: 1767-1773.
- MRC Working Shop, (1971):** Characterization of antisera as reagents. Immunology, 20: 3-10.
- Nadaphan, N.; Wilcox, G. E. and Pexhalf, W. Z. (1982):** Antisera in avian serological assays. Avian Dis., 26: 171-178.
- Narin, R. C. (1969):** Fluorescent protein tracing 3rd Ed. Livingston Edinburgh and London.
- Narin, R. C. and Marrack, J. R. (1964):** Fluorescent protein tracing. 2nd Ed. Livingston Edinburgh and London.
- Noura, F. A.; Albagory, G. F.; Mansour, A. A. and Khodeir, M. H. (2009):** Preparation and evaluation of diagnostic kits of Newcastle disease; infectious bronchitis and infectious bursal disease. Kafrelsheikh Vet. Med. J. 3rd Sci.Cong.46-65.
- Pope, Z. H. and Rowe, W. P. (1964):** Viral genetic material. J. Exp. Med., 120: 121- 125.
- Spendlove, R. S.; Lennette, E. H. and Zohn, A.C. (1963):** Determination of viral antigens. J. Immunol., 90: 554-560.
- Soliman, A. K.; Botros, B. A. M.; Rsiazek, T. G.; Hoogstreal, H.; Helmy, I. and Morin, A. (1989):** Sero-prevalence of Rickettsia typhosa and Rickettsia cornil infection among rodents and dogs in Egypt. Egypt. J. Trop. Med. Hyg. 92: 345-349.
- Tevehia, S. S.; Katz, M. and Rapp, F. (1965):** New surface antigen in cells transformed by Simian Papovirus SV40. Proc.Soc.Exp.Biol. (NY), 119: 896-901.
- Tizard, I.R. (1996):** Veterinary immunology 5th Ed. Saunders Comp., Philadelphia.
- Tumpey, T. M.; Alvarez, R.; Swayme, D. E. and Suarez, D. L. (2005):** A diagnostic aid for differentiating infected from vaccinated poultry based on antibodies to the nonstructural protein (NSP) of influenza A virus. J. Clin. Microbiol. 43: 676-683.
- Watson, B. K. and Coons, A. I. (1954):** Studies of influenza virus infection in the chick embryo using fluorescent antibodies. J. Exp. Med. 99: 419-428.
- Weichselbaum, T. E. (1946):** An accurate and rapid method for the detection of proteins in small amounts of blood serum and plasma. Am. J.Clin.Path.16: 40-49.

دراسات مقارنة على مضادات أمصال الطيور المحملة بالفلوريسين أيزوثيوسينات

تم خلال هذا العمل تحضير أمصال مضادة لأجناس الدجاج والرومي والبط والأوز والحمام والسمان ثم تم ترسيب الجلوبيولين منها وإقرانه بمادة الفلوريسين أيزوثيوسينات. وقد تمت معايرة هذه المقترنات بإجراء اختبار الوميض الفلوريسنتي الغير مباشر على أنتيجينات للنيوكاسل والجمبورو والإلتهاب الكبدي الوبائي في البط وطاعون البط وباراميكسو الحمام باستخدام أجسام مناعية مضادة لكل من هذه الأنتيجينات محضرة نوعياً في طيور محصنة وذلك مقارنة بمقترن مرجعي مضاد للدجاج ١:١٠٠٠٠٠، ١:١٠٠٠٠٠، ١:١٠٠٠٠، ١:١٠٠٠٠، ١:١٠٠٠٠٠، ١:١٠٠٠٠٠٠، ١:١٠٠٠٠٠٠٠، وعند إجراء اختبار الوميض الفلوريسنتي الغير مباشر باستخدام المقترنات المحضرة نوعياً وتبادلياً وجد أن استخدام المقترن النوعي يعطي تفاعل قوي (+٤) بينما الاستخدام التبادلي لمثل هذه المقترنات يعطي تفاعلاً أقل (٣+ إلى ±)، وعلى ذلك يمكن القول بأنه يمكن استخدام أمصال الطيور المختلفة المقترنة بالفلوريسين أيزوثيوسينات في اختبار الوميض الفلوريسنتي المناعي الغير مباشر تبادلياً ولكن بصورة محدودة والأفضل أن يستخدم مضاد جنس الطائر نوعياً للحصول على نتائج أفضل وأكثر دقة.